



Starmaster™ 16% Starter/ Grower Textured Goat



Complete, textured feed designed to be fed to goats from start to finish.

Contains 16% protein	Important for efficient muscle growth.
Contains Decoquinatate	Aids in prevention of Coccidiosis (see tag)
Highly fortified	Promotes optimum performance.
Highly palatable with cooked, steam-flaked grains	Improves palatability, energy digestions, coat, and body condition.
Nutritionally balanced and fortified with vitamins and minerals.	Promotes proper growth, development, reproduction and overall health.
Contains probiotics	Enhances digestion and overall performance.
Contains ammonium chloride.	To acidify the urine, to help avoid "water belly"

DESCRIPTION: Starmaster™ 16% Starter/Grower Textured Goat feed is designed to be fed to weaned goats up to show weight. This high performance feed is formulated to achieve lean muscle growth, optimum body condition, muscle tone, and freshness in goats. Provide clean, fresh water at all times.

FORM

Textured

BAG SIZE

50#



16% Starter/Grower

MEDICATED TEXTURED FEED FOR SHOW GOATS

FOR THE PREVENTION OF COCCIDIOSIS IN YOUNG GOATS CAUSED BY *EIMERIA CHRISTENSENI*
AND *EIMERIA NINAKOLYAKIMOVAE*

ACTIVE DRUG INGREDIENT

DECOQUINATE 22.7 GRAMS/TON

GUARANTEED ANALYSIS

Crude Protein Min 16.00 %
(THIS INCLUDES NOT MORE THAN 2.00% EQUIVALENT CRUDE PROTEIN FROM NON-PROTEIN NITROGEN)
Crude Fat Min 3.50 %
Crude Fiber Max 9.00 %
Calcium Min 0.50 % Max 1.00 %
Phosphorus Min 0.50 %
Salt Min 0.25 % Max 0.75 %
Copper Min 30 ppm Max 35 ppm
Selenium Min 0.30 ppm
Vitamin A Min 7,000 IU/lb

INGREDIENTS

CRACKED CORN, OATS, SOYBEAN MEAL, HEAT PROCESSED SOYBEANS, DISTILLERS DRIED GRAINS WITH SOLUBLES, CANE MOLASSES, VEGETABLE OIL, DEHYDRATED ALFALFA MEAL, COTTONSEED HULLS, WHEAT MIDLINGS, AMMONIUM CHLORIDE, CALCIUM CARBONATE, DICALCIUM PHOSPHATE, SALT, MAGNESIUM OXIDE, SULFUR, FERROUS CARBONATE, FERROUS SULFATE, COPPER SULFATE, MANGANOUS OXIDE, MANGANESE SULFATE, ZINC OXIDE, ZINC SULFATE, COBALT CARBONATE, CALCIUM IODATE, SODIUM SELENITE, VITAMIN A SUPPLEMENT, VITAMIN D₃ SUPPLEMENT, VITAMIN E SUPPLEMENT, SACCHAROMYCES CEREVISIAE, DRIED ENTEROCOCCUS FAECIUM FERMENTATION PRODUCT, DRIED LACTOBACILLUS CASEI, DRIED LACTOBACILLUS ACIDOPHILUS FERMENTATION PRODUCT, DRIED LACTOBACILLUS PLANTARUM FERMENTATION PRODUCT, DRIED LACTOBACILLUS SUBTILIS FERMENTATION PRODUCT, DRIED ASPERGILLUS ORYZAE FERMENTATION EXTRACT, NATURAL FLAVORS.

Feeding Directions: 16% Starter/Grower Goat feed contains 11.35 milligrams of Decoquinatone per pound. Feed 1 pound per 50 pounds of bodyweight to provide 22.7 mg of Decoquinatone per 100 lb of bodyweight. Feed at least 28 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard. Feed with appropriate amounts of roughage to goats. Always provide plenty of clean, fresh water.

Caution: Do not use in feeds containing bentonite.

Warning: Do not feed to goats producing milk for food.

Reminder: Feed is perishable. Store this product in a cool, dry area away from rodents and insects.

Warning: Do not offer any feed that is spoiled, moldy, rodent or insect-infested, or abnormal in appearance or odor, as it may cause illness or death.

Warning: This product contains supplemental copper. Do not feed to sheep or other copper-sensitive species.

7963